

Dermestes frischii* and *D. undulatus* (Coleoptera: Dermestidae) on a Human Corpse in Southern Italy: First Report

Teresa Bonacci^{1,*}, Vannio Vercillo², Mark Benecke³

Abstract: Dermestidae and other necrophagous beetles may aid in the estimation of post-mortem interval, especially in situations where decomposition is advanced and therefore, insect succession patterns are the best method available. We report, for the first time, *Dermestes frischii* Kugelann, 1792 and *D. undulatus* larvae and adults from a human body discovered in Cassano, Cosenza province, southern Italy. The corpse was a 88 year old male. The cadaver was dry and in outdoor location. The other insect species associated with *Dermestes* spp were *Necrobia ruficollis* (adults and larvae) and *Chrysomya albiceps* (Diptera, Calliphoridae) (empty puparium). The time of death was estimated to have been 25 days prior to the finding of the body, based on police investigations and statements of the caretaker. Beetles collected from corpses and crime scenes can be helpful to estimate the colonization interval in summer season and at dry environmental conditions that are to be expected to increase due to climate change.

Key Words: Forensic entomology, *Dermestes frischii*, *Dermestes undulatus*, larder beetles, mummification, Italy, *Chrysomya albiceps*, *Necrobia ruficollis*, time since death.

INTRODUCTION

Larder beetles (Dermestidae) are scavengers that feed on dried tissue and skin of many animals but also on dry tissue of plants. Members of the genus *Dermestes* spp. are distributed all over the planet and can be used in forensic entomology cases (Smith, 1986; Byrd and Castner, 2009; Charabidze *et al.*, 2013; Benecke 2015). Like or together with other Dermestidae, they are indicators of the time since or the season of death (Catts and Goff, 1992), especially when mummified or skeletonized corpses are discovered (Voigt, 1965; Schroeder *et al.*, 2002). Often, just one species of larder beetle is reported on corpses or carcasses (Kumara *et al.*, 2009; Bonacci *et al.*, 2011) but in France in outdoor environments, the two species *D. frischii* and

D. undulatus were found associated (Charabidze *et al.*, 2013). Also, *Necrobia* spp (Coleoptera, Cleridae) was found on human bodies. *D. frischii* is dominant in Iran and reported from animal carcasses in different regions (Kökdener and Polat, 2014; Özdemir and Sert, 2009). *D. frischii* and *D. undulatus* species are active during warm months of the year. Larder beetles are selective in their choice of feeding and breeding substrates, besides avoiding light and excessive humidity (Charabidze *et al.* 2013, Rakowski, 1988, Coombs, 1979; Woodcock *et al.*, 2013). Data on the preferential feeding areas of Dermestidae on the cadaveres were also reported by Voigt (1965) and Schroeder *et al.* (2002). The areas that dry more rapidly than other parts of the body (hands, face and feet) are preferred feeding sites of Dermestids (Charabidze *et al.* 2013).

* In memory of Prof. Dan Dermengiu. He was editor of this journal, director of the "Mina Minovici" National Institute of Legal Medicine in Bucharest, a good friend, and a man wiser and wittier than many. MB

1) University of Calabria, DiBEST Department, Rende, Italy

* Corresponding author: E-mail: teresa.bonacci@unical.it

2) Azienda Sanitaria Provinciale, Sezione di Medicina legale di Cosenza, Italy

3) International Forensic Research & Consulting, Cologne, Germany

CASE DESCRIPTION AND FINDINGS

A 88 years old man was found dead in Cassano allo Jonio, Cosenza province in southern Italy on July, 31 2016. The cause of death was hypertensive myocarditis. The time of death was estimated to have been 22-25 days before the discovery of the body, based on police investigations and testimonies. The man was last seen alive in July, 6 2016. The corpse was found near, 800 meter away from the nursing home where the elderly had been living. The body was found in a supine position dressed in a shirt, pants and socks (Fig. 1) under a *Pistacia lentiscus* bush. The shoes were found close to the body. Autopsy was carried out on August 1, 2016 in Cassano Jonio (CS).

At medico-legal examination, the corpse was found to be in advanced state of decay (mummified stage). The skin was fully dried and of dark-brown color (Fig. 2).

Entomological evidence was collected from and around the body, and from the inside of the bag used to transport the corpse to the autopsy room (Haskell and Catts, 1990; Byrd and Castner, 2009). Live insects (adults and larvae, empty puparia of *Ch. albiceps* and larval exuviae of *Dermestes* spp.) were collected from the corpse during autopsy. Macroscopic medical investigations showed superficial skin lesions on various parts of the body caused by Calliphorid (blowfly) larvae as well as Dermestid and Clerid beetles. The skull was undamaged (Fig 2). No signs of trauma against the osseous structure were observed. The skin and the osseous structure of the thorax was undamaged, and no feeding activity of Calliphorid larvae and Dermestids was observed in these regions.

Numerous larder and Clerid beetles (larvae and adults) were found on the corpse and transferred to the



Figure 1. The cadaver was found in the morning of July, 31 2016, under a *Pistacia lentiscus* bush (A), at 800 meter to the nursing home where the elderly man was hosted (B).



Figure 2. Skin lesions caused by blowfly larvae and Dermestid and Clerid beetles on (A) mummified head (note that the hair is still present) and (B) dried out hands.

laboratory of Entomology of the DiBEST Department, University of Calabria. Particularly, from the pelvis and the legs, many empty larval skins as well as live adults and larvae were found. Empty blowfly puparia (Calliphoridae) were found and collected, mostly from the head and the neck area (Fig. 3), and from the pelvis.

Live Dermestid larvae were reared in laboratory in a plastic container in a climate chamber at $25 \pm 0,5^\circ \text{C}$. The emerging beetles were identified and double-checked by taxonomic keys (Hava and Kadej, 2006; Marcin Kadej, personal information). Dried frass (peritrophic membranes) was found on the corpse, mostly near the genital and pelvic areas (Figs 3 and 4) and on the legs (Fig. 5) where skid lesions had been caused by Dermestid beetles and Calliphorid maggots (Fig. 4).

DISCUSSION

Dermestida are distributed worldwide; they feed on various types of dried animal tissue. The genus *Dermestes* is of forensic importance because it is attracted to human corpses during dry and skeletal stages of decomposition (Pettigrew 1834; Smith 1986; Benecke, 2008; Byrd and Castner, 2009). These beetles are part of the sarcosaprophagous insect succession of the human corpse and animal carcasses (Schroeder *et al.*, 2002; Bonacci *et al.*, 2011; Charabidze *et al.*, 2013; Keshavarzi *et al.*, 2015) in many geographical areas. Usually, the presence of dermestid beetles and their peritrophic membranes on a corpse proves that considerable time has elapsed since the death of the subject (Byrd and Castner, 2009; Benecke, 2015).



Figure 3. Head and neck area (A) where several empty puparium of *Ch. albiceps* were found. Many empty larval skins, and numerous life adults and larvae of dermestid beetles were observed at the pelvis (B).



Figure 4. Peritrophic membranes of *Dermestes* spp., at genital and pelvic areas (A). The dried frass it is frequently found when dermestid beetles have fed on human corpse (B) (Byrd and Castner, 2009; Benecke, 2015).




Figure 5. The dried frass (A) and feeding damage (maybe the result of a mixed activity) by dermestid beetles and maggots (A and B).

Dermestes frischii it is a abundant species found outdoors (Charabidze *et al.*, 2013) and collected in the warm months from human corpses (Kökdener and Polat, 2014) and on animal carcasses (Bana *et al.*, 2012). *D. undulatus* it is very common outside but was never observed in indoor environments. Our findings concerning dermestid beetles are in agreement with previously reported data (Charabidze *et al.*, 2013). In fact, both species (*D. frischii* and *D. undulatus*) were found associated in a dry and hot outdoor environment of southern Italy but, instead, in our reported case, the preferred feeding areas were legs and pelvis. These differences could be explained by different rates of mummification, and by different soft tissue thickness.

Empty puparial shells of *Chrysomya albiceps* suggest that maggots feed rapidly on soft tissue of the head, neck and genital area. It is known that fly larvae need humidity for their development and, a mummified body in a dry and warm habitat it is not a suitable substrate for a long maggots feeding activity. In conclusion, hot and dry environment contributes to rapid mummification

which is a favourable condition for larger beetle activity (feeding and breeding) on the corpse.

Lower as well as very high temperatures and an arid environment disallow the development of Diptera larvae. In such cases, the decay of corpses and carcasses is caused by the activity of other insects like Silphid beetles (Bonacci *et al.*, 2011b). The findings reported here show that *D. frischii* and *D. undulatus* are active during warm months in Calabria and by studying their rates of development in lab conditions and at different experimental temperatures, both species could be used for estimation of post-mortem interval. *Dermestes* species may be found more frequently since due to climate change, warmer and more arid environmental conditions might spread (Lu 2007).

Conflict of interest. The authors declare that there is no conflict of interest.

Acknowledgement. We would like to thank Marcin Kadej for his time and help for the confirmation in the species identification.

References

1. Bana R, Beyarsalan A. Determination of coleoptera species of pig carcasses and internal organs of bovine in Edirne city of Turkey. Bitlis Eren Üniversitesi Fen Bilimleri Dergisi 2012;1:122-126.
2. Benecke M. A brief survey of the history of forensic entomology. Acta Biologica Benrodis 2008;14:15-38.
3. Benecke M. Späte Leichenveränderungen: Besiedlung durch Gliedertiere. In: Brinkmann, B., Madea, B. (eds.) Handbuch gerichtliche Medizin, 2004:Vol. 1, p. 170—187.

4. Benecke M. Arthropods on mummies in the Catacombe dei Cappuccini in Palermo, Italy. 8th International Congress of Dipterology, Potsdam, 2014, Abstract Volume 1, p. 36.
5. Benecke M. So arbeitet die moderne Kriminalbiologie. Cologne: Lübbe, 2015, 10th ed..
6. Bonacci T, Zetto Brandmayr T, Brandmayr P, Vercillo V, Porcelli F. Successional patterns of the insect fauna on a pig carcass in southern Italy and the role of *Crematogaster scutellaris* (Hymenoptera, Formicidae) as a carrion invader. *Entomological Science* 2011;14:125–132.
7. Bonacci T, Greco S, Zetto Brandmayr T. Insect fauna degradation activity of *Thanatophilus* species on carrion in southern Italy (Coleoptera: Silphidae). *Entomol Gener* 2011;33:171-177.
8. Byrd J, Castner J. Forensic entomology: The utility of arthropods in legal investigations. Boca Raton: CRC Press, 2009, 2nd ed..
9. Catts EP, Goff ML. Forensic entomology in criminal investigations. *Annual Review of Entomology* 1992;37:253-272.
10. Charabidze D, Colar T, Vincent B, Pasquerault T, Hedouin V. Involvement of larder beetles (Coleoptera: Dermestidae) on human cadavers: a review of 81 forensic cases. *International Journal of Legal Medicine* 2014;128:1021-1030.
11. Coombs CW. The effect of temperature and humidity upon the development and fecundity fo *Dermestes haemorrhoidalis* Kuster and *Dermestes peruvianus* Laporte de Castelnau (Coleoptera: Dermestidae). *J Stored Prod Res* 1979;15:43-52.
12. Haskell NH, Catts EP. *Entomology and Death: A procedural guide*, Joyce's Print Shop, Clemson, SC, 1990, p. 52-97.
13. Hava J, Kadej M. New data on the occurrence of Dermestidae (Coleoptera) from Hispaniola. Part 1. *Dermestes* Genus. *Polish Journal of Entomology* 2006; 73: 375-377.
14. Keshavarzi D, Moemenbellah-Fard MD, Fereidooni M, Montazeri M. First Report of *Dermestes frischii* Kugelann (Coleoptera: Dermestidae) on a Human Corpse, South of Iran. *Int J Forensic Sci Pathol.* 2015;3: 113-115.
15. Kökdener M, Polat E. Insect succession on dog (*Canis lupus familiaris* L.) carcasses in Samsun province, Turkey. *Munis Entomology & Zoology* 2014;9:858-869.
16. Lu J, Vecchi G, Reichler T. Expansion of the Hadley cell under global warming. *Geophysical Research Letters* 2007;34:L06805.
17. Özdemir S., Sert O. Determination of coleoptera fauna on carcasses in Ankara Provinc, Turkey. *Forensic Science International* 2009;183:24-32.
18. Pettigrew PJ. A history of Egyptian mummies, and an account of the worship and embalming of the sacred animals by the Egyptians; with remarks on the funeral ceremonies of different nations, and observations on the mummies of the Canary islands, of the ancient Peruvians, Burman priests, &c. London: Longman, Reese *et al.*, 1834
19. Rakowski G. Effect of illumination intensity on the response of the hide beetle, *Dermestes maculatus*, to aggregation pheromone. *J Insect Physiol* 1988; 34:1101-1104.
20. Schroeder H, Klotzbach H, Oesterhelweg L, Püschel K. Larder beetles (Coleoptera, Dermestidae) as an accelerating factor for decomposition of a human corpse. *Forensic Sci Int* 2002;127: 231-236.
21. Smith KGV. A manual of forensic entomology. London: Trustees of the British Museum (Natural History), 1986.
22. Voigt J. Specific post-mortem changes produced by larder beetles. *J Forensic Med* 1965;12:76-80.
23. Woodcock L, Gennard D, Eady P. Egg laying preferences and larval performance in *Dermestes maculatus*. *Entomol Exp Appl.* 2013.